



## Parasitology is a tool for identifying the original biotope of the gibel carp (*Carassius auratus gibelio* Berg, 1932)

### Parazitológiai bizonyítékok az ezüstkárász (*Carassius auratus gibelio* Berg, 1932) eredetéről

Molnár K.<sup>1</sup>, Nyeste K.<sup>2</sup>, Székely Cs.<sup>1</sup>

<sup>1</sup>MTA ATK, Állatorvos-tudományi Intézet, Budapest

<sup>2</sup>Debreceni Egyetem TTK, Hidrobiológiai Tanszék, Debrecen

**Keywords:** original biotope of *Carassius* spp., gibel carp, myxosporean infection, nomenclature

**Kulcsszavak:** kárász eredeti biotópja, ezüstkárász, nyálkaspórás fertőzőttség, nevezéktani problémák

#### Abstract

At this time the occurrence of three *Carassius* taxa (*C. carassius*, *C. auratus auratus* and *C. auratus gibelio*) are known from Europe. Crucian carp [*Carassius carassius* (Linnaeus, 1758)] is a native fish species in European waters. The goldfish, a species of Chinese origin arrived to Europe long time ago, and at the time when Linnaeus in 1758 published his *Systema Naturae* he described two *Carassius* species, the crucian carp as *Cyprinus carassius* and the goldfish as *Cyprinus auratus*. During the last two centuries 13 other *Carassius* spp. were described which proved to be synonymous of *C. carassius* and 3-3 species as synonymous of *Carassius auratus auratus* and *C. auratus gibelio*, respectively. The authors confute the European origin of *Carassius gibelio* Bloch, called as Prussian carp. They compared infections of the gibel carp and goldfish with myxosporeans in Europe and in the Far-East and found that these fishes in the Far-East have been infected by several host specific *Myxobolus* and *Thelohanellus* species, while in Europe of them only a single species is known. Great differences in the range of myxosporean spp. suggest that both gibel carp and goldfish are Far-East origin fishes which arrived to Europe in the historical times. The identical myxosporean fauna of the gibel carp (*Carassius auratus gibelio*) and goldfish (*Carassius auratus auratus*) also shows that they are subspecies of *Carassius* (*Cyprinus auratus*) described by Linnaeus.

#### Kivonat

Európában jelenleg három *Carassius* taxa (*C. carassius*, *C. auratus auratus* and *C. auratus gibelio*) előfordulása ismert. Közülük a széles kárász [*Carassius carassius* (Linnaeus, 1758)] mint Európában őshonos halfaj ismert. Az aranyhal egy kínai eredetű halfaj néhány évszázada érkezett Európába, és amikor Linnaeus 1758-ban publikálta a *Systema Naturae* művét abban két *Carassius* faj, a széles kárász és az aranyhal előfordulását jegyezte fel *Cyprinus carassius*, illetve *Cyprinus auratus* néven. Az utóbbi két évszázadban további 13 *Carassius* fajt ítak le, melyek később a *C. carassius* szinonimáinak bizonyultak. További 3-3 faj a *Carassius auratus auratus* ill. az ezüstkárász *C. auratus gibelio* szinonimájának bizonyult. A szerzők a porosz ponty (*Carassius gibelio* Bloch) európai eredetét a kárász fajok távol-keleti és európai nyálkaspórás-parazitafaunájának összevetésével cáfolják. A Távol-Keleten az ezüst kárászt és az aranyhalat számos *Myxobolus* és *Thelohanellus* faj fertőzi, Európában ezek közül csak egyetlen faj ismert. A nyálkaspórás fajok arányában lévő különbségek arra utalnak, hogy az ezüstkárász és aranyhal távol-keleti eredetűek és Európába emberi segítséggel a történelmi időkben érkeztek. Az aranyhal és az ezüstkárász azonos parazitája arra utal, hogy az ezüstkárász (*Carassius auratus gibelio*), valamint az aranyhal (*Carassius auratus auratus*) Linnaeus által leírt *Carassius* (*Cyprinus*) *auratus* alfajai.

## Introduction

Gibel carp, this commonly occurring fish in the Far-East freshwaters, is an aggressively invasive fish species in the Danube region since the second half of the twentieth century. Its European origin is a vexed question among ichthyologists. Some of the specialists (e.g. Kottelat & Freyhof 2007, Froese & Pauli 2018) still accepts that the species is identical with that one described in Prussia by Bloch (1782) as *Cyprinus gibelio*, and they suggest of designating the fish as Prussian carp [*Carassius gibelio* (Bloch, 1782)]. At the other side some authors (e.g. Berg 1932, 1964, Hensel 1971) think that the fish is of Far-East origin, and it is a subspecies of the goldfish [*Carassius auratus* (Linnaeus, 1758)]. They prefer naming the fish as gibel carp, allogynogenetic crucian carp or silver crucian carp (*Carassius auratus gibelio* Berg, 1932). Their view seems to be strongly supported by Kalous et al. (2004), who after re-examining the type specimen of *Carassius gibelio* (Bloch, 1782) in the Natural History Museum in Berlin found that the type specimen was a *Carassius carassius* (Linnaeus, 1758). Hensel (1971) thought that the fish described as gibel carp were in most case goldfish specimens run wild from aquaria. Lelek (1987) supposed that the gibel carp could be accidentally got to Europe with the goldfish but he have not even precluded, that the fish was accurately populated to European waters. Pintér (1980) reported that the gibel carp was first introduced to Hungary in 1954 from Bulgaria. Present authors think that the *Carassius* sp. described by Bloch (1782) as gibel carp (*Cyprinus gibelio*) was a malformed morphological variation of the crucian carp, therefore it is a synonym of *Carassius carassius*. Summary of synonyms of taxa of the genus *Carassius* are presented in *Table 1*. The fish species which at this time is called as gibel carp was introduced to Europe only in the last century. The gibel carp, this commonly occurring fish in the Far-East freshwaters, is an aggressively invasive fish species in the Danube region living here since the second half of the twentieth century. During its expansion gibel carp forced the native crucian carp into isolated refuges in its newly conquered territories. The fact that gibel carp this invasive fish did not spread fast from Prussia to other parts of Europe and even this time it is not inhabitant of that region contradicts to its existence in Bloch's ages. Though these facts seem unanimously prove the Asian origin of the gibel carp, present authors by comparing the parasite fauna of the European and Asian population of the gibel carp presents further evidences in this matter. A similar work was published recently on the Asian origin of the common carp by Molnár (2009), who comparing the parasitic infection of the European common carp (*Cyprinus carpio* Linnaeus 1758) and the Asian common carp provided similar data.

Examination of the parasite fauna is a useful tool for identification of the origin of small fish groups or a whole population in Oceans (MacKenzie, 2002). By this method spooning places of Pacific Ocean fishes in small rivers of Alaska or Kamchatka could exactly be outlined (Arthur & Albert 1993, Konovalov 1995, Timi, 2007). In this respect host specific parasites, may play a specific role, due to the fact that in general these introduced fishes in the new habitat harbour less number of specific parasites than in the original biotope (Torchin et al. 2003). Sagarin et al. 2006) described, that according to the parasite/predator escape hypothesis, an introduced species may profit from the favorable situation in the new habitat, and attains higher population densities and greater individual sizes in the colonized areas compared to the conspecifics in their native range. On the other hand, Prenter et al. (2004) thought that introduced hosts may bring with them all or a subset of their native parasite fauna into the new area or they may even adopt local parasites. The importance of parasites has also been recognized in invasions when native host populations are infected by a new parasite transported with introduced host (Vooren 1972, Moravec & Taraschewski 1988). In most case the advantage of a new species at a new habitat comes from the fact that the introduced species left the majority of its specific parasites and pathogens in the native habitat.

Most of the parasites can be used as tools, but species with strict host specificity or species infecting only some closely related fishes are the best for this purpose. Among these

parasites the host specific species of Monogenea and Myxosporea deserve a special attention.

Table 1. Synonyms of taxa of the genus *Carassius*  
1. táblázat. A *Carassius nembe tartozó taxonok szinonimái*

Taxon	Author and date/Leíró és dátum
<b><i>Carassius carassius</i></b>	<b>(Linnaeus, 1858)</b>
<i>C. auratus wui</i>	Tchang, 1930
<i>C. carax</i>	(Lesniewski, 1837)
<i>C. coeruleus</i>	Basilewski, 1855
<i>C. discolor</i>	Basilewski, 1855
<i>C. gibelio minutes</i>	(Kessler, 1856)
<i>C. humilis</i>	Heckel, 1837
<i>C. limnaei</i>	Bonaparte, 1845
<i>C. limnei</i>	Malm, 1877
<i>C. moles</i>	Nordmann, 1840
<i>C. oblongus</i>	Hecket et Kner, 1858
<i>C. pekingensis</i>	Basilewsky, 1855
<i>C. vulgaris</i>	Nordmann, 1840
<i>C. vulgaris subventrosus</i>	Walecki, 1863
<b><i>Carassius auratus auratus</i></b>	<b>(Linnaeus, 1758)</b>
<i>C. auratus cantonensis</i>	Tchang, 1933
<i>C. chinensis</i>	Gronow, 1854
<i>C. encobia</i>	Bonaparte, 1845
<b><i>Carassius auratus gibelio</i></b>	<b>Berg 1932</b>
<i>C. ellipticus</i>	Heckel, 1848
<i>C. vulgaris kolenti</i>	Gronow, 1854
<i>C. vulgaris ventrosus</i>	Walecki, 1863

In this paper, besides historical data and field observations by analysing the myxosporidian infections of Asian and European gibel carp parasitological evidences are presented which show that the parasite fauna of the Far East gibel carp stock is significantly richer in species than that of the European gibel carp. The parasitic infection of the gibel carp in Europe harbours only a small part of species infecting this fish in Chinese waters.

### Material and methods

A systematic research on the parasite fauna of fishes including *Carassius* spp. started in Hungary in 1960. Since this time the parasitic infection of freshwater and pond-cultured fishes has been regularly surveyed, and all parasitic groups infecting fishes have been concerned. During these surveys complete parasitological dissections were performed; histological sections from the collected material were prepared; samples from parasites were preserved in alcohol or as slide preparations; photos and video images were recorded. A special attention was paid to parasites of invasive fishes (gibel carp, brown bullhead [(*Ameiurus nebulosus* (Lesueur, 1819))], pumpkinseed [(*Lepomis gibbosus* (Linnaeus, 1758))], Ponto-Caspian gobies), the colonisation of which in the Hungarian fauna took place in the last century, or to parasites infecting fishes which were regularly introduced during these ages such as European eel [*Anguilla anguilla* (Linnaeus, 1758)], goldfish, koi-carp, silver carp [*Hypophthalmichthys molitrix* (Valenciennes, 1844)], bighead carp [*Hypophthalmichthys nobilis* (Richardson, 1845)] and grass carp [*Ctenopharyngodon idella* (Valenciennes, 1844)].

Data to the present paper have been collected from the records of the authors since 1960. In the first years of investigations rivers and lakes between the Danube and Tisza and the Pannonian region of Hungary were populated only with the crucian carp. In 1960 to 1964 32 specimens of 7 to 20 cm long crucian carp were studied for parasitic infection.

Investigation of the parasite fauna of the gibel carp started in 1964 in the ponds and dead arms of the Hármas-Körös River in Szarvas (East Hungary, where some years before this fish species was introduced from Bulgaria). From 1964 to 1984 parasites of 42 specimens of 3 to 26 cm long gibel carp were studied in this habitat and up to 2006 further 21 specimens were examined from different territories of Hungary. Parasitic infection of the goldfish was studied in the Temperate Water Fish Hatchery, in Százhalombatta (near to Budapest), where the goldfish was cultured in ponds. Of this fish species 44 specimens of 3 to 15 cm long goldfish were examined from 1987 to 2006. In 2007 10 specimens of crucian carp collected from isolated ponds (refuges) of south-west Hungary were also examined for parasites.

### Results and discussion

Some fish species as the pumpkinseed, brown bullhead arriving from North America became the member of the European fish fauna due to unconsidered fish introductions more than 150 years ago. These fishes enriched the parasite fauna in Europe with some monogeneans as *Cleidodiscus pricei* Mueller, 1936 or *Haplocladius dispar* (Mueller, 1936) and *H. similis* (Mueller, 1936) (Roman 1960). The expansion of the Ponto-Caspian gobies takes places in these days conquering the whole Danube region and inhabiting even the northern part of the American continent. Their expansion is partially an active invasion but ballast water of ships contributes to their rapid spread as well (Ahnelt et al. 1989; Jude et al. 1992). Of the specific parasites these fishes introduced only some coccidian species to the Hungarian stretches of the Danube River from the Black Sea biotope (Molnár 2006).

Eels arrived to Hungary exclusively by anthropogenic transfer, and since 1961 to 1991 in each year glass eels were introduced into Lake Balaton and some closed system water basins. Until 1990 the parasite fauna was the same as in natural ways populated rivers and lakes in Europe (Murai 1971). In the first part of the 1980<sup>th</sup> years an invasive and pathogenic nematode, *Anguillicola crassus* Kuwahara, Niimi & Itagaki, 1974 arrived to Europe due to the introduction of the Japanese eel, (*Anguilla japonica* Temminck & Schlegel, 1846), and this parasite infected the European eel and caused a heavy infection in this fish (Hartman 1987). This parasite was first detected in Hungary in 1990 (Székely et al. 1991). Another large-scale anthropochore fish transfer took place in the years of 1950<sup>th</sup> when the Amur River fishes were introduced to the European part of the former Soviet Union. At this time stocks of the Amur wild carp (*Cyprinus carpio erythropterus*), grasscarp [*Ctenopharyngodon idellus* (Valenciennes)], silver carp [*Hypophthalmichthys molitrix* (Valenciennes)] and bighead carp [*Hypophthalmichthys nobilis* (Richardson)] arrived regularly to Europe. These latter fishes introduced several pathogenic parasites to the European continent (Musselius 1967) and most of the parasites became introduced also to Hungary (Szakolczai & Molnár, 1963). This was also the period when the colour variation of the Japanese carp (koi) arrived to the Western part of Europe more frequently using the fastest way of fish transfer, the airway routs. In most cases fishes, among them herbivorous fishes introduced their specific parasites to the new biotope, where they were able to survive on their original hosts. Some of them, however, introduced some less specific parasites, as well, which were infective also to the native fishes. Of the latter the cestode *Bothriocephalus acheilognathi* Yamaguti, 1934 is the best known, which became a world-wide pathogen (Malewitskaya 1958).

Of the introduced parasites, species of three fishes, *Cyprinus carpio haematopterus*, *Carassius auratus auratus* and *Carassius auratus gibelio*, deserve a special attention. These fishes after the anthropogenic breakdown of biogeographic barriers got in contact with endemic subspecies and species of their genera (*Cyprinus carpio carpio*, *Carassius carassius*) in the new areas, and the non-native and native species could infect each other with their specific parasites.

Table 2. *Myxosporeans described from taxa of the genus Carassius*  
2. táblázat. A *Carassius nembe tartozó taxonokból leírt nyálkaspóráások*

Table 2a. *Myxobolus spp. described from crucian carp*  
2a. táblázat. Széles kárászból leírt *Myxobolus* fajok

Name of the parasite/Parazitafaj neve	Infected organs/ Fertőzött szerv	Fish species as described/Gazdaállat	Locality/Helyszín
<i>Myxobolus carassii</i> Klokachewa, 1914	liver, intest., abdominal cav.	<i>Carassius carassius</i>	Russia
<i>M. diversus</i> Nie & Li, 1973	gills, kidney	<i>Carassius carassius cuvieri</i>	China
<i>M. elongatus</i> Fujita, 1924	kidney	<i>Carassius carassius</i>	Japan
<i>M. thelohanellus</i> Shulman & Vikhrova, 1952	gills	<i>Carassius carassius</i>	Russia

Table 2b. *Myxobolus spp. described from goldfish*  
2b. táblázat. Aranyhalból leírt *Myxobolus* fajok

Name of the parasite/Parazitafaj neve	Infected organs/ Fertőzött szerv	Fish species as described/Gazdaállat	Locality/Helyszín
<i>M. bladderia</i> Chen & Ma, 1998	gall-bladder	<i>Carassius auratus auratus</i>	China
<i>M. cantonensis</i> Chen, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>M. cultus</i> Yokoyama, Ogawa & Wakabayashi, 1995	cartilage	<i>Carassius auratus</i>	Japan
<i>M. echengensis</i> Chen, 1998	kidney	<i>Carassius auratus auratus</i>	China
<i>M. egregius</i> Li & Nie, 1973	gills, kidney, spleen	<i>Carassius auratus auratus</i>	China
<i>M. hearti</i> Chen, 1998	heart	<i>Carassius auratus auratus</i>	China
<i>M. hokiangensis</i> Ma, 1998	ureter, urinary bladder	<i>Carassius auratus auratus</i>	China
<i>M. huananensis</i> Chen, 1998	skin, gills	<i>Carassius auratus auratus</i>	China
<i>M. huchowensis</i> Chen, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>M. kingchowensis</i> Ma & Chen, 1998	almost all organs	<i>Carassius auratus auratus</i>	China
<i>M. lokiaensis</i> Chen, 1998	urinary bladder	<i>Carassius auratus auratus</i>	China
<i>M. nanyangensis</i> (Hu, 1965) Eiras et al., 2005	gills	<i>Carassius auratus auratus</i>	China
<i>M. nanyuensis</i> Chen, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>M. pekingensis</i> Chen, 1998	intestine, liver, kidney	<i>Carassius auratus auratus</i>	China
<i>M. pseudosquaræ</i> Chen, 1998	gills, caudal fin	<i>Carassius auratus auratus</i>	China
<i>M. pyramidis</i> Chen, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>M. qingyiensis</i> ( <i>Myxobolus yaanensis</i> Ma & Zhao, 1998) Eiras et al, 1965 syn.	spleen, abdominal cavity	<i>Carassius auratus auratus</i>	China
<i>M. tachengensis</i> Chen, 1998	gall-bladder, front intestine	<i>Carassius auratus auratus</i>	China
<i>M. tuberculosis</i> Nie & Li, 1992	heart, urinary bladder, ureter	<i>Carassius auratus auratus</i>	China
<i>M. tunghuensis</i> Chen, 1998	urinary bladder, kidney	<i>Carassius auratus auratus</i>	China
<i>M. urinarybladderi</i> ( <i>Myxosoma tunghuensis</i> Chen, 1998) Eiras et al., 2005	urinary bladder	<i>Carassius auratus auratus</i>	China
<i>M. valatus</i> Li & Nie, 1973	gills, intestine, kidney, skin	<i>Carassius auratus auratus</i>	China
<i>M. wasjugani</i> Bocharova & Donec, 1974	muscles	<i>Carassius auratus</i>	River Ob
<i>M. wuhanensis</i> Chen, 1998	kidney, gall-bladder	<i>Carassius auratus auratus</i>	China
<i>M. wulii</i> ( <i>Myxosoma magna</i> Wu & Li, 1986) Landsberg & Lom, 1991	gills, spleen, abdominal cavity	<i>Carassius auratus auratus</i>	China
<i>M. wushingensis</i> Chen, 1998	kidney, intestine	<i>Carassius auratus auratus</i>	China

For studying differences in the parasite fauna of the original habitat and the new biotope two parasitic groups are the best stools, namely monogeneans and myxosporeans. Monogeneans, which develop by a direct way from fish to fish, and could be transmitted from continents to continents even by a single fish hosts, might arrived from the Far East to Europe as early as the first Portuguese sailors introduced the goldfish into Europe. Therefore, in the present paper their role has not been considered. Studying the myxosporean infection of fishes might serve more information in this respect. Myxosporeans, among them *Myxobolus* spp. develop by oligochaete alternative hosts and their colonization at a new habitat is more difficult. When comparing the *Myxobolus* fauna of the Far East *Carassius* spp. with those of the European stocks, a huge difference can be seen. From the crucian carp only 4 *Myxobolus* spp have been described (Table 2a), and only 2 of them (*M. carassii* and *M. thelohanellus*) are known in Europe. At the other side from the goldfish 26 *Myxobolus* spp, and from the gibel carp 13 *Myxobolus* spp. have been described

from the waters of China, Japan and the Amur Basin. (Table 2b,c). In Hungary only a single species (*M. carassii*) is known from the crucian carp, and another species, *M. diversus* from the goldfish (Molnár & Székely 2003). This species was later on recorded on the fins of the gibel carp, as well (Székely & Molnár unpublished data). Studying differences in parasitic infections of *Carassius* spp. in the Far East and Europe another myxosporean group *Thelohanellus* gives convincing picture (Table 2d). From the Far East *Carassius* spp. 8 *Thelohanellus* spp. have been described, from Europe neither of them is known. Up to this time the occurrence of five myxosporean species is known from Hungary (Table 2 e).

The large number of *Myxobolus* spp. infecting the Far-East *Carassius* spp. and their scarce number in Europe suggest that the original habitat of the goldfish and the gibel carp was the Chinese region. In case of the goldfish it is an accepted view, in case of gibel carp, however, the Far-East origin is disputed. The fact, that a *Carassius* species was described in Europe as *Carassius gibelio* by Bloch in 1782 (Bloch 1782) seemed to support the European origin of this species. Considering, however, that the specimen described by Bloch (Bloch 1782) was indeed a crucian carp indicate that the homeland of the gibel carp should be looked for in the Chinese-Amur fauna region. Besides our data, received from studying the myxosporean fauna of *Carassius* spp, historical data prove, that the gibel carp arrived to the Carpathian Basin rather late. Knowing the aggressive invading nature of this fish one cannot accept that this fish was an original member of the European fish fauna. The gibel carp forced back the wide ranged crucian carp into remote refugees within some decades after arriving to the Danube system. No similar action was recorded in the past Prussian territories.

Table 2c. *Myxobolus* spp. described from gibel carp  
2c. táblázat. Ezüstkárászról leírt *Myxobolus* fajok

Name of the parasite/Parazitafaj neve	Infected organs/ Fertőzött szerv	Fish species as described/Gazdaállat	Locality/Helyszín
<i>M. acutus</i> ( <i>Sphaerospora acuta</i> Fujita, 1912) Landsberg & Lom, 1991	gills	<i>Carassius auratus gibelio</i>	Japan
<i>M. alacaudatus</i> Yukhimenko, 1986	gills, muscles, fins	<i>Carassius auratus gibelio</i>	Amur basin
<i>M. artus</i> Akhmerov, 1960	kidney	<i>Carassius auratus gibelio</i>	Amur basin
<i>M. gibelio</i> Yukhimenko, 1986	gills, fins, kidney	<i>Carassius auratus gibelio</i>	Amur basin
<i>M. gibelioi</i> Wu & Wang, 1982	gills	<i>Carassius auratus gibelio</i>	China
<i>M. honghuensis</i> Liu, Whipps, Gu & Huang, 2012	pharynx	<i>Carassius auratus gibelio</i>	China
<i>M. kubanicus</i> Bykhovskaya-Pavlovskaya & Bykhovski, 1940	gut, muscles	<i>Carassius auratus gibelio</i>	Kuban River
<i>M. lentisuturalis</i> Dyková, Fiala & Nie, 2002	muscle fibres	<i>Carassius gibelio</i>	China
<i>M. orientalis</i> Shulman, 1962	gills	<i>Carassius auratus gibelio</i>	China
<i>M. platyrostris</i> Akhmerov, 1960	kidney	<i>Carassius auratus gibelio</i>	Amur basin
<i>M. pseudoparvus</i> Li & Nie, 1973	skin	<i>Carassius auratus gibelio</i>	China
<i>M. sacchalinesis</i> ( <i>Lentospora sacchalinesis</i> Fujita, 1923) Landsberg & Lom, 1991	kidney	<i>Carassius auratus gibelio</i>	Japan
<i>M. solidus</i> Shulman, 1962	gills	<i>Carassius auratus gibelio</i>	China
<i>M. sphaericus</i> ( <i>Lentospora spherica</i> Fujita, 1924) Landsberg & Lom, 1991	kidney	<i>Carassius auratus gibelio</i>	Japan
<i>M. turpisrotundus</i> Zhang, Wang, Li, & Gong, (2010)	skin, fins	<i>Carassius auratus gibelio</i>	China

Table 2d. *Thelohanellus* spp. described from taxa of the genus *Carassius*  
2d. táblázat. A *Carassius* nembe tartozó taxonokból leírt *Thelohanellus* fajok

Name of the parasite/Parazitafaj neve	Infected organs/ Fertőzött szerv	Fish species as described/Gazdaállat	Locality/Helyszín
<i>T. carassii</i> Akhmerov, 1960	gills	<i>Carassius auratus gibelio</i>	Amur River, Far-East Russia
<i>T. hupehensis</i> Ni et Li, 1992	spleen, kidney	<i>Carassius auratus auratus</i>	China
<i>T. liaohoensis</i> Chen in Chen et Ma, 1998	gall bladder	<i>Carassius auratus auratus</i>	China
<i>T. membranicaudata</i> Sun, 2006	gills	<i>Carassius auratus auratus</i>	China
<i>T. nanhaiensis</i> Chen in Chen et Ma, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>T. oliviformis</i> Wu et Wang, 1982	muscle	<i>Carassius auratus gibelio</i>	China
<i>T. relortus</i> Chen in Chen et Ma, 1998	gills	<i>Carassius auratus auratus</i>	China
<i>T. wangi</i> Zhang, 2013	gills	<i>Carassius auratus gibelio</i>	China
<i>T. wuhanensis</i> Xiao et Chen, 1993	skin	<i>Carassius auratus gibelio</i>	China

Table 2e. *Myxosporeans found in taxa of the genus Carassius in Hungary*  
 2e. táblázat. A *Carassius nembe tartozó taxonokból megtalált nyálkaspóráások Magyarországon*

Name of the parasite/ Parazitafaj neve	Infected organs/ Fertőzött szerv	Fish species as described/Gazdaállat	Described in/ Eredeti leírási helyszín
<i>Myxobolus carassii</i> Klokachewa, 1914	liver, intest., abdominal cav.	<i>Carassius carassius</i>	Russia
<i>M. diversus</i> Nie & Li, 1973	fins	<i>Carassius auratus</i>	China
<i>Sphaerospora carassii</i> Kudo, 1919	gills	<i>Carassius gibelio</i> , <i>C. auratus</i>	Japan
<i>Hoferellus carassii</i> Achmerov, 1960	kidney	<i>Carassius gibelio</i>	Amur River
<i>Myxobilatus</i> sp.	kidney	<i>Carassius auratus</i> , <i>Carassius gibelio</i>	
<i>Myxidium</i> sp.	kidney	<i>Carassius auratus</i> , <i>Carassius gibelio</i>	

Molecular biological studies indicate that DNA sequences of the goldfish and the gibel carp resemble to each other much better than to the crucian carp (Jia et al. 2006; Zhu et al. 2006; Sakai et al. 2009).

It comes from the uncertain origin of the *Carassius* spp. that there is a great confusion both in the Latin and English names. This situation characterizes first of all Chinese papers. No matter the two species, *Carassius carassius* and *C. auratus* described by Linnaeus (Linnaeus 1758), have commonly accepted names as crucian carp and goldfish, respectively. Unfortunately in the Chinese literature *C. auratus* is often called as crucian carp, red crucian carp, etc. The problem is more complex at the gibel carp. The common parasite fauna and molecular evidences prove that the goldfish and the gibel carp belong to one species *Carassius auratus* (Linnaeus, 1758) which has two subspecies *Carassius auratus auratus* (Linnaeus, 1758) and *Carassius auratus gibelio* Berg, 1932. The name *Carassius gibelio* (Bloch, 1782) should be regarded as a junior synonym of *C. carassius* (Linnaeus, 1758). At a similar way designation of the species to Prussian carp is wrong, as the species has nothing to do with the Past Prussia (Molnár 2015).

#### Acknowledgement

The research was supported by MAHOP-2.1.1-2016-2017-00002. (A horgászati- és halgazdálkodás szempontból jelentős halfajok tenyésztését és termelését támogató technológia-, tudástranszfer és innovációs infrastruktúra fejlesztése)

The research was financed by the Higher Education Institutional Excellence Programme of the Ministry of Human Capacities in Hungary, within the framework of the 4th thematic programme of the University of Debrecen. Krisztián Nyeste was supported by ÚNKP-18-3 New National Excellence Program of the Ministry of Human Capacities.

#### References

- Arthur, J. R., Albert, E. (1993): Use of parasites for separating stocks of greenland halibut (*Reinhardtius hypoglossoides*) in the Canadian Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 2175–2181.
- Ahnelt, H. (1989): Die Marmorierte Grundel (*Proterorhinus marmoratus* (Pallas), Pisces: Gobiidae)—ein postglazialer Einwanderer. *Österreichs Fischerei* 42: 11–14.
- Arthur, J. R., Albert, E. (1993): Use of parasites for separating stocks of greenland halibut (*Reinhardtius hypoglossoides*) in the Canadian Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 2175–2181.
- Bloch, M.E. (1782): Oeconomische Naturgeschichte der Fische Deutschlands. In: Bloch, M.E.: *Allgemeine Naturgeschichte der Fische*, Berlin, pp. 226..
- Berg, L. S. (1932): Über *Carassius carassius* und *C. gibelio*. *Zoologischer Anzeiger* 98: 15–18.
- Berg, L. S. (1964): *Freshwater fishes of the USSR and adjacent countries*, Vol. 2. Israel Program for Scientific Translations, Jerusalem, pp. 496.
- Froese, R., Pauly, D. (Eds.) (2018): FishBase. World Wide Web electronic publication. www.fishbase.org, version (26/07/2018).
- Hensel, K. (1971): Some notes on the systematic status of *Carassius auratus gibelio* (Bloch, 1782) with further record of this fish from the Danube river in Czechoslovakia. *Vestník Československe Spolecnosti Zoologicke* 35/3: 186–198.
- Jia, Z. Y., Sun, X. W., Liang, L. Q., Lu, C. Y., Lei, Q. Q. (2006): Isolation and characterization of microsatellite markers from Fangzheng silver crucian carp, *Carassius auratus gibelio* (Bloch), and cross-amplification in

- the closely related species crucian carp, *Carassius auratus auratus* (Linnaeus). *Molecular Ecology Notes* 6:1141–1143
- Jude, D. J., Reider, R. H., Smith, G. R. (1992): Establishment of Gobiidae in the Great-Lakes Basin. *Canadian Journal of Fisheries and Aquatic Sciences* 49: 416–421.
- Kalous, L., Bohlen, J., Ráb, P. (2004): What fish is *Carassius gibelio*: Taxonomic and nomenclatoric notes. *Proceedings of XI. European Congress of Ichthyology*. Tallinn, Estonia, September 6-10, 2004, Book of abstracts 26–27.
- Konovalov, S. M. (1995): Parasites as indicators of biological processes, with special reference to sockeye salmon (*Oncorhynchus nerca*). *Canadian Journal of Fisheries and Aquatic Sciences* 52/S1: 202–212.
- Kottelat, M., Freyhof, J. (2007): *Handbook of European freshwater fishes*. Publications Kottelat, Cornol, Switzerland, pp. 646.
- Lelek, A. (1987): Threatened Fishes of Europe. p. 171-172. In: . *The Freshwater Fishes of Europe*. Vol. 9. AULA-Verlag, Wiesbaden.
- Linnaeus, C. (1758). *Systema naturae*. 10th edition, Vol. 1. Laurentii Salvii, Stockholm, pp. 824.
- MacKenzie, K. (2002): Parasites as biological tags in population studies of marine organisms: an update. *Parasitology* 124/S: 153–163.
- Molnár K. (2006): Some remarks on parasitic infections on the invasive *Neogobius* spp. (Pisces) in the Hungarian reaches of the Danube River, with description of *Goussia szekelyi* sp. n. (Apicomplexa: Eimeriidae). *Journal of Applied Ichthyology* 22: 395–400.
- Molnár K. (2009): Data on the parasite fauna of the European common carp *Cyprinus carpio* and Asian common carp *Cyprinus carpio haematopterus* support an Asian Ancestry of the species. *Aquaculture, Aquarium, Conservation & Legislation International Journal of the Bioflux Society* 2: 391–400.
- Molnár K. (2015): Néhány szó az ezüst kárászról, eredetéről, behurcolásáról, betegségeiről és jelentőségéről. *Halászat* 108/1: 26–27
- Molnár K., Székely Cs. (2003): Infection in the fins of the goldfish *Carasius auratus* caused by *Myxobolus diversus* (Myxosporea). *Folia Parasitologica* 50: 31–36.
- Murai É. (1971): Tapeworms (Cestodes) parasitizing Eels introduced into Lake Balaton. *Parasitologia Hungarica* 4: 145–156.
- Musselius, V. A. (1967): *Parasites and diseases of phytophagous fishes and prevention*. KOLOS, Moscow, pp. 93 (in Russian).
- Pintér K (1980): Exotic fishes in Hungarian waters: their importance in fishery utilization of natural water bodies and fish farming. *Fisheries Management* 11/4: 163–167.
- Prenter J., MacNeil C., Dick J. T. A., Dunn A. M. (2004): Roles of parasites in animal invasions. *Trends in Ecology & Evolution* 19/7: 385–390.
- Roman, E. (1960): Plathelminthes, Class Monogenoidea. In: Botnariuc, N. (ed.). *Fauna of the Romanian People's Republic*, vol. 2. Editura Academiei Române, Bucharest, pp. 149.
- Sagarin, R. D., Gaines, S. D., Gaylord, B. (2006): Moving beyond assumptions to understand abundance distributions across the ranges of species. *Trends in Ecology & Evolution* 21: 524–530.
- Sakai, H., Iguchi, K., Yamazaki, Y., Sideleva, V. G., Goto, A. (2009): Morphological and mtDNA sequence studies on three crucian carps (*Carassius*: Cyprinidae) including a new stock from the Ob River system, Kazakhstan. *Journal of Fish Biology* 74: 1756–1773.
- Székely Cs., Láng M., Csaba Gy. (1991): First occurrence of *Anguillicola crassus* in Hungary. *Bulletin of the European Association of Fish Pathologists* 11: 162–163.
- Szakolczai J., Molnár K. (1966): Veterinärmedizinische Untersuchungen an den in Ungarn eingebürgerten pflanzenfressenden Fischarten. *Z. F. Fischerei NFBD*. 14: 139–152.
- Timi, J. T. (2007): A multiple scale look at parasites as biological tags in marine fish: from host stocks to regional fish assemblages. Abstracts of 7th International Symposium on Fish Parasites. *Parassitologia* 49/S2: 252.
- Torchin, M. E., Lafferty, K. B., Dobson, A. P., McKenzie, V. J., Kuris, A. M. (2003): Introduced species and their missing parasites. *Nature* 421: 628–630.
- Vooren, C. M. (1972): Ecological aspects of the introduction of fish species into natural habitats in Europe, with special reference to the Netherlands. *Journal of Fish Biology* 4: 565–583.
- Zhu, H.P., Ma, D.M., Gui, J. F. (2006): Triploid origin of the gibel carp as revealed by 5S rDNA localization and chromosome painting. *Chromosome Research* 14:767–776.

**Authors:**

Kálmán MOLNÁR (molnar.kalman@agrar.mta.hu), Krisztián NYESTE (nyeste.krisztian@science.unideb.hu),  
CSABA SZÉKELY (szekely.csaba@agrar.mta.hu)